

WEAK WEIGHTED INEQUALITIES FOR A DYADIC ONE-SIDED MAXIMAL FUNCTION IN $\mathbb{R}^n$

SHELDY OMBROSI

ABSTRACT. In this note we introduce a dyadic one-sided maximal function defined as $M^{+,d}f(x) = \sup_{Q \text{ dyadic}: x \in Q} \frac{1}{|Q|} \int_{Q^+} |f|$, where Q^+ is a certain cube associated with the dyadic cube Q and $f \in L^1_{loc}(\mathbb{R}^n)$. We characterize the pair of weights (w, v) for which the maximal operator $M^{+,d}$ applies $L^p(v)$ into weak- $L^p(w)$ for $1 \leq p < \infty$.

1. INTRODUCTION, NOTATION AND MAIN RESULTS

In [10] E. Sawyer introduced the class of weights A_p^+ , which characterizes the pair of weights (w, v) such that the one-sided Hardy-Littlewood maximal operator apply $L^p(v)$ into weak- $L^p(w)$. These classes of weights and their associated theory have been the subject of much study. Different proofs of Sawyer's results and more general extensions are given in the papers of F. J. Martín-Reyes, Ortega-Salvador and A. de la Torre [6], [4] and [5]. In [1] H. Aimar, L. Forzani and F. J. Martín-Reyes proved the existence of one-sided singular integral operators, and L. de Rosa and C. Segovia have developed great part of the theory of one-sided weighted Hardy spaces (see [8] and [9]). However, as far as the author knows, the theory of one-sided weights has been developed **only** in $\mathbb{R}$.

The results in this note try to give a first step to overcome that limitation. We will define a dyadic one-sided maximal function in $\mathbb{R}^n$ and we characterize the pair of weights (w, v) such that the operator associated with this maximal function applies $L^p(v)$ into weak- $L^p(w)$ (Theorem 1.2 below).

If $I = [a, b]$ is a bounded interval we denote $I^+ = [b, 2b - a]$ and $I^- = [2a - b, a]$. If $Q = I_1 \times I_2 \times \dots \times I_n$ is a cube in $\mathbb{R}^n$, we denote $Q^+ = I_1^+ \times I_2^+ \times \dots \times I_n^+$, and $Q^- = I_1^- \times I_2^- \times \dots \times I_n^-$.

Given $f \in L^1_{loc}(\mathbb{R}^n)$, we define the dyadic one-sided maximal function $M^{+,d}f(x)$ as

$$M^{+,d}f(x) = \sup_{Q \text{ dyadic}: x \in Q} \frac{1}{|Q|} \int_{Q^+} |f|,$$

In a similar way $M^{-,d}f(x) = \sup_{Q \text{ dyadic}: x \in Q} \frac{1}{|Q|} \int_{Q^-} |f|$. In dimension one, this maximal function and, more generally, dyadic one-sided fractional maximal functions were defined in [5].

Let $x \in \mathbb{R}^n$, $x = (x_1, x_2, \dots, x_n)$, and let h be a positive real number. We denote $Q_{x,h} = [x_1, x_1 + h) \times [x_2, x_2 + h) \times \dots \times [x_n, x_n + h)$, $Q_{x,h-} = [x_1 - h, x_1) \times [x_2 - h, x_2) \times \dots \times [x_n - h, x_n)$, and $Q_{x,h}^{\frac{1}{2}} = [x_1 + \frac{h}{2}, x_1 + h) \times [x_2 + \frac{h}{2}, x_2 + h) \times \dots \times [x_n + \frac{h}{2}, x_n + h)$.

Key words and phrases. **weights, dyadic one-sided maximal function**
2000 *Mathematics Subject Classification* Primary: 42B25; Secondary: 28B99.

With this notation, we define the following maximal functions

$$M^+f(x) = \sup_{h>0} \frac{1}{|Q_{x,h}|} \int_{Q_{x,h}} |f|, \quad M^-f(x) = \sup_{h>0} \frac{1}{|Q_{x,h-}|} \int_{Q_{x,h-}} |f|, \text{ and}$$

$$N^+f(x) = \sup_{h>0} \frac{1}{|Q_{x,h}^{\frac{1}{2}}|} \int_{Q_{x,h}^{\frac{1}{2}}} |f|,$$

In dimension $n = 1$, the maximal functions $M^{+,d}f(x)$, $N^+f(x)$, and $M^+f(x)$ are pointwise equivalents (see [5]), this fact is not true if $n > 1$.

Let $Q = [a_1, b_1] \times [a_2, b_2] \times \dots \times [a_n, b_n]$, with $b_i - a_i = h$, and let s be a positive real number. We denote $(Q)^{s,+} = [a_1, a_1 + sh] \times [a_2, a_2 + sh] \times \dots \times [a_n, a_n + sh]$, and $(Q)^{s,-} = [b_1 - sh, b_1] \times [b_2 - sh, b_2] \times \dots \times [b_n - sh, b_n]$. We observe that $|(Q)^{s,+}| = |(Q)^{s,-}| = s^n |Q|$ and $Q = (Q)^{1,+} = (Q)^{1,-}$.

Remark 1.1. Let R and Q be cubes such that $R^+ \subset Q^+$, then $R \subset (Q)^{2,+}$.

As usual, a weight $w(x)$ is a measurable and non-negative function. If $E \subset \mathbb{R}^n$ is a Lebesgue measurable set, we denote its w -measure by $w(E) = \int_E w(t)dt$.

We will say that a pair of weights (w, v) satisfies the condition $A_p^+(\mathbb{R}^n)$, (or (w, v) belongs to the class $A_p^+(\mathbb{R}^n)$) $p > 1$, if there exists a constant C such that for every cube Q ,

$$(1) \quad \int_Q w \left(\int_{Q^+} v^{-\frac{1}{p-1}} \right)^{p-1} \leq C |Q|^p.$$

The condition $A_1^+(\mathbb{R}^n)$ is given by

$$(2) \quad M^-w(x) \leq Cv(x) \quad \text{a.e.}$$

Interesting gemetric properties of this classes of weights were obtained by Martín-Reyes in [3]

We define similar classes of weights on dyadic cubes. We will say that a pair of weights (w, v) satisfies the condition $A_p^{+,d}(\mathbb{R}^n)$ ($p > 1$) if (1) holds for every dyadic cube Q , and the condition $A_1^{+,d}$ is given by $M^{-,d}w(x) \leq Cv(x)$ a. e.

It is clear that if $w \geq 0$ is a nondecreasing function in each coordinate, the pair (w, w) satisfies the condition $A_p^+(\mathbb{R}^n)$ and therefore $A_p^{+,d}(\mathbb{R}^n)$. Thus, these conditions are weaker than the Muckenhoupt's condition $A_p(\mathbb{R}^n)$ introduced in [7], which characterizes the pair of weights such that the Hardy-Littlewood maximal operator applies $L^p(v)$ into weak- $L^p(w)$.

The next theorem solves a conjecture of A. de la Torre and F. J. Martín-Reyes.

Theorem 1.2. Let w and v be weights on $\mathbb{R}^n$ and let $1 \leq p < \infty$. Then, the following conditions are equivalent:

- (a) The pair of weights (w, v) satisfies the condition $A_p^{+,d}(\mathbb{R}^n)$
- (b) The maximal operator $M^{+,d}$ is of weak type (p, p) with respect to the pair (w, v) , that is: there is a constant C such that for every function $f \in L_{loc}^1(\mathbb{R}^n)$ and every $\lambda > 0$

$$w(\{x : M^{+,d}f(x) > \lambda\}) \leq C\lambda^{-p} \int_{\mathbb{R}^n} |f|^p v$$

Theorem 1.3. *Let w and v be two weights on $\mathbb{R}^n$ and let $1 \leq p < \infty$. If (w, v) belongs to the class $A_p^+(\mathbb{R}^n)$ then the maximal operator N^+f is of weak type (p, p) with respect to the pair (w, v) .*

2. PROOF OF THE RESULTS

Let f be an integrable and non-negative function and let λ be a positive real number. We denote $\Omega_\lambda = \{x : M^{+,d}f(x) > \lambda\}$. It is not hard to see that there exists a disjoint collection of dyadic cubes $\{Q_j\}$ such that

$$\cup_j Q_j = \Omega_\lambda \text{ and for each } j, f_{Q_j^+} = \frac{1}{|Q_j|} \int_{Q_j^+} f > \lambda.$$

In general, the associated collection $\{Q_j^+\}$ is not disjoint. This fact is the main

difficulty that appearing in our proof of Theorem 1.2. However with the next lemma we overcome that difficulty.

Lemma 2.1. *Let $1 \leq p < \infty$. Let (w, v) be a pair of weights satisfying the condition $A_p^{+,d}(\mathbb{R}^n)$. Let μ be a positive real number. If $\{Q_j\}_{j \in \Gamma_\mu}$ is a disjoint collection of dyadic cubes (Γ_μ denotes the family of indexes j) such that for each $j \in \Gamma_\mu$, $\mu < f_{Q_j^+} \leq 2\mu$, then*

$$\sum_{j \in \Gamma_\mu} w(Q_j) \leq \frac{2^{p+n+2}C}{\mu^p} \int_{\cup_{j \in \Gamma_\mu} Q_j^+} |f|^p v,$$

where C is the smallest constant that we can take in the condition $A_p^{+,d}(\mathbb{R}^n)$ for the pair (w, v) .

Proof. For each integer $m \geq 0$, we define the subfamily of indexes i_m in Γ_μ as

$$i_m = \{j \in \Gamma_\mu : \text{there exist exactly } m \text{ cubes } Q_s^+ : Q_j^+ \subsetneq Q_s^+ \text{ with } s \in \Gamma_\mu\},$$

and we denote

$$\sigma_m = \cup_{j \in i_m} Q_j^+.$$

Remarks:

- (1) It is clear that $\Gamma_\mu = \cup_{m \geq 0} i_m$. Furthermore, if j_1, j_2 belong to i_m then $Q_{j_1}^+ \cap Q_{j_2}^+ = \emptyset$, and therefore

$$\sum_{j \in i_m} \int_{Q_j^+} f = \int_{\sigma_m} f.$$

- (2) If $j \in i_{m+1}$ ($m \geq 0$) then for each integer k , $0 \leq k \leq m$, there is only one $j_k \in i_k$ such that $Q_j^+ \subset Q_{j_k}^+$. Thus $\sigma_{m+1} \subset \sigma_m$ and $\int_{\sigma_{m+1}} f \leq \int_{\sigma_m} f$.
- (3) We fix a non negative integer m_0 and $j_0 \in i_{m_0}$. If $Q_j^+ \subsetneq Q_{j_0}^+$ then $j \in i_m$ where $m > m_0$, and moreover, by remark 1.1 we have that $Q_j \subset (Q_{j_0})^{2,+}$.

Taking into account the last observation and since the cubes Q_j are disjoint we have

$$\cup_{m > m_0} \cup_{\substack{j \in i_m : \\ Q_j^+ \subsetneq Q_{j_0}^+}} Q_j \subset (Q_{j_0})^{2,+},$$

and

$$\sum_{m>m_0} \sum_{\substack{j \in i_m: \\ Q_j^+ \subsetneq Q_{j_0}^+}} |Q_j| \leq 2^n |Q_{j_0}|.$$

Therefore, since for every $j \in \Gamma_\mu$, $\mu < f_{Q_j^+} \leq 2\mu$, we get

$$\begin{aligned} \sum_{m>m_0} \int_{\sigma_m \cap Q_{j_0}^+} f &= \sum_{m>m_0} \sum_{\substack{j \in i_m: \\ Q_j^+ \subsetneq Q_{j_0}^+}} \int_{Q_j^+} f \\ &\leq 2\mu \sum_{m>m_0} \sum_{\substack{j \in i_m: \\ Q_j^+ \subsetneq Q_{j_0}^+}} |Q_j| \leq 22^n \mu |Q_{j_0}| < 2^{n+1} \int_{Q_{j_0}^+} f. \end{aligned}$$

Thus, we have that $\sum_{m=m_0+1}^{m_0+2^{n+2}} \int_{\sigma_m \cap Q_{j_0}^+} f < 2^{n+1} \int_{Q_{j_0}^+} f$. Then, there is a index m , $m_0+1 \leq m \leq m_0+2^{n+2}$, such that $\int_{\sigma_m \cap Q_{j_0}^+} f < \frac{1}{2} \int_{Q_{j_0}^+} f$, and since $\int_{\sigma_{m+1} \cap Q_{j_0}^+} f \leq \int_{\sigma_m \cap Q_{j_0}^+} f$, we have that, if $j_0 \in m_0$,

$$\int_{Q_{j_0}^+ \cap \sigma_{m_0+2^{n+2}}} f < \frac{1}{2} \int_{Q_{j_0}^+} f,$$

which implies that

$$\int_{Q_{j_0}^+ - \sigma_{m_0+2^{n+2}}} f > \frac{1}{2} \int_{Q_{j_0}^+} f.$$

From the last inequality and since $\mu < f_{Q_{j_0}^+}$, we get

$$(3) \quad \frac{1}{|Q_{j_0}|} \int_{Q_{j_0}^+ - \sigma_{m_0+2^{n+2}}} f > \frac{\mu}{2}.$$

Now we will estimate $\sum_{j \in \Gamma_\mu} w(Q_j)$. We suppose $p > 1$. From (3), Hölder's inequality, and by condition $A_p^{+,d}(\mathbb{R}^n)$, we have

$$\begin{aligned} \sum_{j \in \Gamma_\mu} w(Q_j) &= \sum_{m=0}^{\infty} \sum_{j \in i_m} w(Q_j) \leq \left(\frac{2}{\mu}\right)^p \sum_{m=0}^{\infty} \sum_{j \in i_m} w(Q_j) \left(\frac{1}{|Q_j|} \int_{Q_j^+ - \sigma_{m+2^{n+2}}} f\right)^p \\ &\leq \left(\frac{2}{\mu}\right)^p \sum_{m=0}^{\infty} \sum_{j \in i_m} w(Q_j) \left[v^{-\frac{1}{p-1}} (Q_j^+)\right]^{p-1} \frac{1}{|Q_j|^p} \int_{Q_j^+ - \sigma_{m+2^{n+2}}} f^p v \\ (4) \quad &\leq \frac{2^p C}{\mu^p} \sum_{m=0}^{\infty} \int_{\sigma_m - \sigma_{m+2^{n+2}}} f^p v. \end{aligned}$$

(If $p = 1$ we simply observe that $\frac{w(Q_j)}{|Q_j|} \int_{Q_j^+ - \sigma_{m+2^{n+2}}} f \leq \int_{Q_j^+ - \sigma_{m+2^{n+2}}} f M^{-,d} w$, and by condition $A_1^{+,d}(\mathbb{R}^n)$ we can obtain the previous estimate also in this case).

Then, by (4) and since

$$\begin{aligned} \sum_{m=0}^{\infty} \int_{\sigma_m - \sigma_{m+2^{n+2}}} f^p v &= \sum_{k=0}^{2^{n+2}-1} \sum_{m=0}^{\infty} \int_{\sigma_{2^{n+2}m+k} - \sigma_{2^{n+2}(m+1)+k}} f^p v \\ &= \sum_{k=0}^{2^{n+2}-1} \int_{\sigma_k} f^p v \leq 2^{n+2} \int_{\sigma_0} f^p v, \end{aligned}$$

we obtain the lemma. ■

Proof of Theorem 1.2 . We will prove that $(a) \Rightarrow (b)$ since $(b) \Rightarrow (a)$ can be obtained, with simple changes, by a standard argument (e.g., see pp. 390 in [2]).

Assume that $(w, v) \in A_p^{+,d}(\mathbb{R}^n)$. We will estimate

$$w(\{x : M^{+,d}f(x) > \lambda\}) = w(\Omega_\lambda).$$

Recall that we can write $\Omega_\lambda = \cup_j Q_j$, where $f_{Q_j^+} > \lambda$.

For each non-negative integer k , we consider the subfamily of indexes $c_k = \{j : 2^k \lambda < f_{Q_j^+} \leq 2^{k+1} \lambda\}$. Then, we can apply Lemma 2.1 to the collection $\{Q_j\}_{j \in c_k}$, with $\mu = 2^k \lambda$, getting

$$\sum_{j \in c_k} w(Q_j) \leq \frac{2^{p+n+2}C}{(2^k \lambda)^p} \int_{\mathbb{R}^n} f^p v.$$

Therefore

$$\begin{aligned} w(\Omega_\lambda) &= \sum_j w(Q_j) = \sum_{k=0}^{\infty} \sum_{j \in c_k} w(Q_j) \\ &\leq \sum_{k=0}^{\infty} \frac{2^{p+n+2}C}{(2^k \lambda)^p} \int_{\mathbb{R}^n} f^p v = \frac{2^{p+n}8C}{\lambda^p} \int_{\mathbb{R}^n} f^p v. \end{aligned}$$

which proves that $(a) \Rightarrow (b)$. ■

Proof of Theorem 1.3 . Fix an integrable and non-negative function f . We know that

$$(5) \quad \{x : M^{+,d}f(x) > 4^{-n}\lambda\} = \cup_j Q_j,$$

where $\{Q_j\}_j$ is a disjoint collection of dyadic cubes $\{Q_j\}_j$ such that for each j , $f_{Q_j^+} > 4^{-n}\lambda$. We will prove that

$$(6) \quad \{x : N^+f(x) > \lambda\} \subset \cup_j (Q_j)^{4,-}.$$

If $x \in \{x : N^+f(x) > \lambda\}$ there exists $h > 0$ such that $f_{Q_{x,h}^{\frac{1}{2}}} > \lambda$. Let k be the

integer such that $2^{k+1} \leq h < 2^{k+2}$. Then, there exist at most 4^n dyadic cubes R such that $R \cap Q_{x,h}^{\frac{1}{2}} \neq \emptyset$ and $|R| = 2^{kn}$, and therefore one of them satisfies that $\int_R f \geq 4^{-n} \int_{Q_{x,h}^r} f$. Then, since $|R| = 2^{kn} \leq |Q_{x,h}^{\frac{1}{2}}|$, we have that

$$\frac{1}{|R|} \int_R f \geq \frac{4^{-n}}{|R|} \int_{Q_{x,h}^r} f \geq \frac{4^{-n}}{|Q_{x,h}^{\frac{1}{2}}|} \int_{Q_{x,h}^{\frac{1}{2}}} f > 4^{-n}\lambda,$$

and therefore the dyadic cube $R^- \subset \{x : M^{+,d}f(x) > 4^{-n}\lambda\}$. Thus there exists a

maximal dyadic cube Q_j in the collection in (5) such that $R^- \subset Q_j$, thus $(R^-)^{4,-} \subset (Q_j)^{4,-}$ and since $x \in (R^-)^{4,-}$ we obtain (6).

Now, by condition $A_p^+(\mathbb{R}^n)$, we have that

$$\begin{aligned} \int_{(Q_j)^{4,-}} w \left(\int_{(Q_j^+)^{4,+}} v^{-\frac{1}{p-1}} \right)^{p-1} &\leq C |Q_j|^p, \text{ if } p > 1, \text{ and} \\ \frac{1}{|Q_j|} \int_{(Q_j)^{4,-}} w &\leq C M^- w(x) \leq C v(x) \text{ for a.e } x \in Q_j^+, \text{ if } p = 1. \end{aligned}$$

Thus, arguing as in the proof of Lemma 2.1 and taking into account the last inequalities, we can prove that

$$\sum_{j: 2^{k-2n}\lambda < f_{Q_j^+} \leq 2^{k-2n+1}\lambda} w((Q_j)^{4,-}) \leq \frac{C}{(2^k\lambda)^p} \int_{\mathbb{R}^n} |f|^p v.$$

Then, following the reasoning as in the proof of Theorem 1.2 we can obtain the theorem. ■

Final Remark

We have already observed that if $n > 1$ the maximal functions $N^+f(x)$ and $M^+f(x)$ are not pointwise equivalents. However, we believe that it is possible to prove that if the pair of weights (w, v) satisfies the condition $A_p^+(\mathbb{R}^n)$ then the

maximal operator M^+ is of weak type (p, p) with respect to (w, v) .

I would like to thank the referee and P. Panzone for their helpful comments.

REFERENCES

- [1] H. Aimar, L. Forzani and F. Martín-Reyes, *On weighted inequalities for singular integrals*. Proc. Amer. Math. Soc. 125, (1995), 2057-2064.
- [2] J. García Cuerva, and J. L. Rubio de Francia, *Weighted norm inequalities and related topics*, North Holland, (1985).
- [3] F. J. Martín-Reyes, *On the one-sided Hardy-Littlewood maximal function in the real line and in dimensions greater than one*. Fourier analysis and partial differential equations (Miraflores de la Sierra, 1992), 237-250, Stud. Adv. Math., CRC, Boca Raton, FL, 1995.
- [4] F. J. Martín-Reyes, *New proofs of weighted inequalities for the one-sided Hardy-Littlewood maximal functions*, Proc. Amer. Math. Soc. 117 (1993), 691-698.
- [5] F. J. Martín-Reyes and A. de la Torre, *Two weight norm inequalities for fractional one-sided maximal operators*, Proc. Amer. Math. Soc. 117 (1992), 691-698.
- [6] F. J. Martín-Reyes, P. Ortega-Salvador, and A. de la Torre, *Weighted inequalities for one-sided maximal functions*. Trans. Amer. Math. Soc. 319 (1990), 517-534.
- [7] B. Muckenhoupt, *Weighted inequalities for the Hardy maximal function*. Trans. Amer. Math. Soc. 165 (1972), 207-226.
- [8] L. de Rosa and C. Segovia, *Weighted H^p spaces for one sided maximal functions*, Contemporary Math. 189, (1995) 161-183.
- [9] L. de Rosa and C. Segovia, *Dual spaces for one-sided weighted Hardy spaces*, Rev. Un. Mat. Arg. (1997).
- [10] E. Sawyer, *Weighted inequalities for the one-sided Hardy-Littlewood maximal functions*, Trans. Amer. Math. Soc. 297 (1986), 53-61.

Sheldy Ombrosi.

mail adress:

Depto. de Matemática, Universidad Nacional del Sur,
Avenida Alem 1253, Bahía Blanca, Buenos Aires, ARGENTINA.
e-mail: sombrosi@uns.edu.ar